

**Reduced Order Model Approach for Imaging with Waves**

**Josselin GARNIER**

**Centre de Mathematiques Appliquees, Ecole Polytechnique, France**

**Email: [josselin.garnier@polytechnique.edu](mailto:josselin.garnier@polytechnique.edu)**

We consider the inverse problem for the scalar wave equation. Sensors probe the unknown medium to be imaged with a pulse and measure the backscattered waves. The objective is to estimate the velocity map from the array response matrix of the sensors. Under such circumstances, conventional Full Waveform Inversion (FWI) can be carried out by nonlinear least-squares data fitting. It turns out that the FWI misfit function is high-dimensional and non-convex and it has many local minima. A novel approach to FWI based on a data driven reduced order model (ROM) of the wave equation operator is introduced and it is shown that the minimization of ROM misfit function performs much better. The talk is based on a joint work with L. Borcea (Univ. Michigan), A. Mamonov (Univ. Houston), J. Zimmerling (Uppsala Univ.).

References:

- [1] L Borcea, J Garnier, AV Mamonov, J Zimmerling, When data driven reduced order modeling meets waveform inversion, arXiv:2302.05988
- [2] L Borcea, J Garnier, AV Mamonov, J Zimmerling, Waveform inversion with a data driven estimate of the internal wave, SIAM Journal on Imaging Sciences 16 (1), 2023, 280-312.
- [3] L Borcea, J Garnier, AV Mamonov, J Zimmerling, Waveform inversion via reduced order modeling, Geophysics 88 (2), 2023, R175-R191.